

**LIZOBON**

Each film coated tablet contains:  
Linezolid IP 600 mg  
Colour : Titanium Dioxide IP

**Dosage :**  
As directed by the Physician.

**Storage :**  
Store protected from light & moisture, at a temperature not exceeding 30°C.

Keep all medicines out of reach of children.

Mfg. Lic. No. : 31/UA/2013

**SCHEDULE H PRESCRIPTION  
DRUG - CAUTION**  
Not to be sold by retail without the prescription of a Registered Medical Practitioner.

Manufactured by :  
Pure & Cure Healthcare Pvt. Ltd.  
(A subsidiary of **Akums Drugs & Pharmaceuticals Ltd.**)  
Plot No. 26A, 27-30, Sector-8A,  
I.I.E., SIDCUL, Ranipur,  
Haridwar-249 403, Uttarakhand

Kindly Scan the QR code  
for more information on  
pack insert



Marketed By :

**BONGYN**  
Bioscience Pvt. Ltd.  
Shed No 2-B, G. F. Block No 896,  
Near Shree Kanta Packaging  
Opp Devam-4 Estate, Santej Super  
City Road Santej-382721  
Taluka: Kalol District: Gandhinagar

**Brand Name:** LIZOBON

**Composition:** Linezolid IP 600 mg Tablets

**General Information:**

LIZOBON is a premium product from **Bongyn Bioscience Pvt Ltd**, designed to effectively treat bacterial infections, particularly those resistant to other antibiotics. Each tablet contains **Linezolid IP 600 mg**, a synthetic antibiotic belonging to the oxazolidinone class. It works by inhibiting bacterial protein synthesis, thereby stopping the growth and spread of bacteria. Manufactured under stringent quality standards, Lizobon ensures efficacy and safety for patients.

**Drug Class:**

- Linezolid: Oxazolidinone antibiotic

**Indications and Uses:**

- Treatment of pneumonia, including community-acquired and hospital-acquired pneumonia
- Skin and soft tissue infections caused by Gram-positive bacteria
- Infections caused by resistant bacteria, including MRSA (Methicillin-resistant Staphylococcus aureus) and VRE (Vancomycin-resistant Enterococcus)
- Complicated and uncomplicated bacterial infections

---

**Common and Less Common Side Effects:****Common Side Effects:**

- Diarrhea
- Nausea
- Headache
- Vomiting
- Changes in taste sensation

**Less Common Side Effects:**

- Peripheral neuropathy
- Low blood counts (thrombocytopenia, anemia)
- Lactic acidosis (rare)
- Skin rash or itching

If any side effects persist or worsen, contact your doctor immediately.

---

**Doses and Administration:**

- **Recommended Dose:** One tablet (600 mg) every 12 hours or as directed by your doctor.
- Swallow the tablet whole with water, with or without food.
- Complete the full course of treatment even if you start feeling better.

---

**Warnings and Precautions:**

- Monitor blood counts regularly during prolonged use to avoid potential hematological effects.
- Inform your doctor if you have a history of anemia, bone marrow suppression, or infections.
- Avoid high tyramine-containing foods (e.g., aged cheese, fermented products) during treatment to prevent hypertensive episodes.
- Use with caution in patients with a history of seizures or neurological disorders.

---

**Drug Interactions:**

- Inform your doctor if you are taking serotonergic drugs (e.g., SSRIs, MAO inhibitors) as these may increase the risk of serotonin syndrome.
- Avoid concurrent use with decongestants, vasopressors, or dopaminergic agents.

- Do not use with other antibiotics unless recommended by your healthcare provider.
- 

**Contraindications:**

- Hypersensitivity to Linezolid or any of the tablet's components
  - Patients taking MAO inhibitors or within two weeks of stopping such treatment
  - Uncontrolled hypertension or hyperthyroidism
- 

**Dos and Don'ts:****Dos:**

- Take Lizobon exactly as prescribed by your doctor.
- Inform your doctor of all medications and supplements you are taking.
- Keep hydrated during the treatment.

**Don'ts:**

- Do not skip doses or stop the medication prematurely.
  - Avoid alcohol during treatment.
  - Do not use this medication without a doctor's prescription.
- 

**Before Taking Lizobon:**

- Discuss your medical history with your doctor, especially if you have liver or kidney issues, hypertension, or a history of seizures.
  - If you are pregnant, planning pregnancy, or breastfeeding, inform your doctor to determine if the medication is suitable for you.
  - Report any known allergies or adverse reactions to antibiotics.
- 

**Contact Information:**

**Bongyn Bioscience Pvt Ltd**

**Address:** 11, First Floor, Science City Road, Sola, Ahmedabad

**Phone:** +91 8160 351 340

**Email:** [info@bongyn.com](mailto:info@bongyn.com)

**Website:** [www.bongyn.com](http://www.bongyn.com)

---

**Disclaimer:**

This is general information about Lizobon. Always use this medication under the guidance of your

doctor. Do not change the prescribed dose or stop taking the medication without consulting your healthcare provider.

**All rights reserved to Bongyn Bioscience Pvt Ltd. 2025**