



Brand Name: LORNIBON-P

Composition: Lornoxicam 8 mg + Paracetamol IP 325 mg Tablets

General Information:

LORNIBON-P is a premium product from **Bongyn Bioscience Pvt Ltd**, designed to provide effective relief from pain and inflammation. Each tablet contains **Lornoxicam 8 mg**, a non-steroidal anti-inflammatory drug (NSAID), and **Paracetamol 325 mg**, an antipyretic and analgesic. This combination works synergistically to reduce pain, inflammation, and fever. Manufactured under stringent quality standards, LORNIBON-P ensures optimal patient care and safety.

Drug Class:

- Lornoxicam: Non-Steroidal Anti-Inflammatory Drug (NSAID)
- Paracetamol: Antipyretic and Analgesic

Indications and Uses:

- Management of acute and chronic pain conditions, such as osteoarthritis and rheumatoid arthritis
- Relief from musculoskeletal and joint pain
- Alleviation of moderate to severe pain associated with inflammation
- Reduction of fever and associated discomfort

Common and Less Common Side Effects:

Common Side Effects:

- Nausea
- Vomiting
- Stomach pain
- Diarrhea
- Dizziness

Less Common Side Effects:

- Skin rash
- Constipation
- Headache
- Increased liver enzymes
- Allergic reactions

If any side effect persists or worsens, seek medical attention promptly.

Doses and Administration:

- **Recommended Dose:** One tablet once or twice daily, as advised by your doctor.
- Swallow the tablet whole with water, preferably after meals to reduce gastrointestinal discomfort.
- Do not exceed the prescribed dose.

Warnings and Precautions:

- Use with caution in patients with a history of gastrointestinal ulcers, bleeding, or perforations.
- Regular monitoring is advised for patients with liver or kidney impairments.
- Avoid using LORNIBON-P during the last trimester of pregnancy.
- Patients with cardiovascular conditions should consult their doctor before use.
- Prolonged use may require monitoring for liver function.

Drug Interactions:

- Avoid concurrent use with other NSAIDs or corticosteroids to reduce the risk of gastrointestinal side effects.
 - May interact with anticoagulants, increasing the risk of bleeding.
 - Diuretics and ACE inhibitors may reduce the effectiveness of LORNIBON-P.
 - Inform your doctor about all medications and supplements you are taking.
-

Contraindications:

- Hypersensitivity to Lornoxicam, Paracetamol, or any excipient in the tablet
 - Patients with severe liver or kidney impairment
 - Active or history of peptic ulcer disease or gastrointestinal bleeding
 - Severe heart failure
-

Dos and Don'ts:**Dos:**

- Take LORNIBON-P exactly as prescribed by your doctor.
- Inform your doctor of any pre-existing conditions or ongoing treatments.
- Stay hydrated and maintain a balanced diet during the treatment.

Don'ts:

- Do not self-medicate or adjust the dose without consulting your doctor.
 - Avoid alcohol consumption during treatment.
 - Do not use this medication if you are pregnant or breastfeeding without medical advice.
-

Before Taking LORNIBON-P:

- Inform your doctor about any allergies or hypersensitivities.
 - Discuss your medical history, especially if you have liver, kidney, or heart conditions.
 - Notify your doctor if you are pregnant, planning to conceive, or breastfeeding.
-

Contact Information:

Bongyn Bioscience Pvt Ltd

Address: 11, First Floor, Science City Road, Sola, Ahmedabad

Phone: +91 8160 351 340

Email: info@bongyn.com

Website: www.bongyn.com

Disclaimer:

This is general information about LORNIBON-P. Always use this medication under the guidance of your doctor. Do not change or discontinue the prescribed dose without medical consultation.

All rights reserved to Bongyn Bioscience Pvt Ltd. 2025